

VU Research Portal

A Taylor series expansion approach to the functional approximation of finite queues

Abbas, K.; Heidergott, B.F.; Aissani, D.

2011

document version

Early version, also known as pre-print

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Abbas, K., Heidergott, B. F., & Aissani, D. (2011). *A Taylor series expansion approach to the functional approximation of finite queues*. (Research Memorandum; No. 2011-49). Faculty of Economics and Business Administration.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

A Taylor series expansion approach to the functional approximation of finite queues

Research Memorandum 2011-49

**Karim Abbas
Bernd Heidergott
Djamil Aïssani**

A Taylor Series Expansion Approach to the Functional Approximation of Finite Queues

Karim Abbas*, Bernd Heidergott†, Djamil Aïssani‡

Abstract

This paper presents a new approach to the functional approximation of the M/G/1/N built on a Taylor series approach. Specifically, we establish an approximative expression for the remainder term of the Taylor series that can be computed in an efficient manner. As we will illustrate with numerical examples, the resulting Taylor series approximation turns out to be of practical value.

keywords: Series expansion approach; Taylor series; $M/G/1/N$ queue; Performance Measures; Deviation matrix

1 Introduction

Queueing models are a well-established tool for the analysis of stochastic systems from areas as diverse as manufacturing, telecommunication, transport and the service industry. Typically, a queueing model is a simplified representation of the real-world system under consideration. In addition, often there is not sufficient statistical data to determine the service and interarrival time distribution, or, in case the type of distribution is known, there is statical uncertainty on the exact values of the parameters of the distribution. For these reasons, perturbation analysis of queueing systems (PAQS) has been developed. PAQS studies the dependence of the performance of a given queueing system on the underlying distributional assumptions. Consider, for example, an M/M/1 model with interarrival rate λ and service rate μ and suppose that one is interested in the variance of the stationary queue length. A first question addressed by PAQS is that on what the effect of small change in, say, λ on the performance would be. This type of perturbation analysis is called *parameter sensitivity*. In addition, PAQS addresses the issue what would happen if, say, the service time distribution is uniform on a certain interval rather than exponential. This

*LAMOS, University of Bejaia, Algeria

†Dept. Econometrics, Vrije Universiteit Amsterdam, the Netherlands

‡LAMOS, University of Bejaia, Algeria

is called *distributional sensitivity*. With the advent of sample path techniques, such as infinitesimal perturbation analysis [10, 5, 2], the score function method [18, 20], and measure-valued differentiation [17], parameter sensitivity is well understood.

The distributional sensitivity is less developed. Importance sampling, see e.g. [19], is a sample path method applicable to this problem. Unfortunately, importance sampling suffers from large variance if the nominal and the perturbed distribution are not ‘sufficiently close’ and it requires that the perturbed distribution is absolutely continuous with respect to the nominal one. For these reasons, most authors use numerical analysis to come up with bounds on the effect of a distributional perturbation. See, for example, the strong stability method [12, 16].

In this paper we present a new approach to PAQS. Firstly, we comprise parameter and distributional sensitivity in one framework. This is achieved by interpreting the distributional sensitivity as a parametric problem: we assume that the distribution under consideration is a convex combination of the nominal and the perturbed distribution, where the weight factor of the convex combination introduces an artificial parameter into the model. Secondly, we develop the performance of the system under consideration into a Taylor series with respect to the parameter of interest, where we make use of a fundamental result on Taylor series for Markov chains, see [7].

For explicatory purposes, we will consider in the following the single server queue with Poisson arrival process and generally distributed service times. Service following the first-come first-served (FCFS) discipline, and there is a limited capacity of in total N customers that can be present at the queue (the one in service included). Customers that do not find an empty place at the queue upon their arrival are lost. Let π^* denote the stationary distribution of the continuous-time queue-length process of the M/G/1N queue. It is well known that π^* can be expressed via the stationary distribution of the Markov chain embedded at departure points of customers, denoted by π . Specifically, let ρ denote the traffic rate, then it holds that

$$\pi^*(i) = \frac{\pi(i)}{\pi(0) + \rho}, i = 0, \dots, N - 1, \quad (1)$$

and

$$\pi^*(N) = \frac{1}{\rho} \left(\rho - 1 + \frac{\pi(0)}{\pi(0) + \rho} \right) = \frac{\pi(0) + \rho - 1}{\pi(0) + \rho}, \quad (2)$$

see [6] for details.

For the PAQS presented in this paper we consider π^* as a mapping of some real-valued parameter θ , in notation π_θ^* . For example, θ may denote the mean service time of the queue. We are interested in obtaining the *functional* dependence of $\pi^*(\theta)$ on θ in a simplified form. For our approach we will compute π_θ^* for some parameter value θ numerically. However, then we will approximate the function $\pi^*(\theta + \Delta)$ on some Δ -interval. More specifically, we will approximate

$\pi^*(\theta + \Delta)$ by a polynomial in Δ . To achieve this we will use the Taylor series expansion approach established in [7]. More specifically, let π_θ denote the stationary distribution of the queue-length process embedded at departure epochs in the M/G/1/N queue, where $\theta \in \mathbb{R}$ denotes a control parameter. Under quite general conditions it holds that $\pi_{\theta+\Delta}$ can be developed into a Taylor series of the following form

$$\pi_{\theta+\Delta} = \sum_{n=0}^k \frac{\Delta^n}{n!} \pi_\theta^{(n)},$$

where $\pi_\theta^{(n)}$ denotes the n -th order derivative of π_θ with respect to θ . We call

$$H_\theta(k, \Delta) = \sum_{n=0}^k \frac{\Delta^n}{n!} \pi_\theta^{(n)}$$

the k -th order Taylor approximation of $\pi_{\theta+\Delta}$ at θ , and

$$r_\theta(k, \Delta) = \pi_{\theta+\Delta} - H_\theta(k, \Delta)$$

the k -th order remainder term at θ . Let η denote the width of the interval $(\theta - \eta, \theta + \eta)$ on which the stationary distribution has to be approximated.

The usefulness of any Taylor series based approach relies on two factors:

1. Fast convergence of the series (a Taylor polynomial of small order yields already a satisfying approximation), i.e., if for small k it holds that

$$\sup_{|\Delta| \leq \eta} |r_\theta(k, \Delta)| \tag{3}$$

is sufficiently small.

2. The ability of computing the remainder term of the Taylor series in an efficient way so that order of the Taylor polynomial that is sufficient to achieve a desired precision of the approximation can be decided *a priori*.

The contribution of the paper is as follows. We investigate the applicability use of the Taylor polynomial $H_\theta(k, \Delta)$ for numerical purposes for M/G/1/N queue. Specifically,

- we identify a recursive form of the derivatives which simplifies coding the algorithm;
- our numerical studies show that already a Taylor series of small order yields good approximations (this addresses topic (1) above);
- a simplified and easily computable expression bounding the remainder of the Taylor series, see (3), is established (this addresses topic (2) above).

The paper is organized as follows. The embedded Markov chain of the $M/G/1/N$ model is presented in Section 2. Our series expansions approach is detailed in Section 3. Numerical examples are provided in Section 4.

We conclude the introduction with a brief discussion of implications of our approach to the numerical approximation of the $M/G/1/\infty$ queue. The stationary distribution π of the embedded jump chain can be obtained in a closed-form only for special cases. This has lead to a rich literature on approximations of the stationary distribution π^* . A very efficient numerical approach is to approximate the $M/G/1/N$ queue by the $M/G/1/\infty$ queue, see [24, 23]. However, this approach is limited to the cases where the traffic load is less than one (so that the corresponding $M/G/1/\infty$ model is stable). For a detailed overview on numerical approaches to the $M/G/1/N$ queue we refer to the excellent survey in [22]. The approach presented in this paper yields a new numerical approach to the $M/G/1/N$ queue that is also feasible for traffic loads larger than one. In addition, our approach yields an approximation of a performance functional on an entire interval and allows for an error bound of the approximation that holds uniformly on an interval.

2 The $M/G/1/N$ Queue

Consider the $M/G/1/N$ queue, where customers arrive according to a Poisson process with rate λ and demand an independent and identically distributed service time with common distribution function $B(t)$ with mean $1/\mu$. There can at most be N customers be present at the queue (including the one in service), and customers attempting to enter the queue when there are already N customers present are lost. The service discipline is FCFS.

Let $X(t)$ denote the number of customers in the $M/G/1/N$ queue at time t , for $t \geq 0$. Note that the queue-length processes $\{X(t) : t \geq 0\}$ of the $M/G/1/N$ system fails to be a Markov process because the service time distribution does not have the memoryless property. Since we have assumed that customers that do not find an empty buffer place upon their arrival are lost, the stationary distribution of $\{X(t) : t \geq 0\}$, denoted by π , exists (independent of the traffic rate). Let $\{X_n : n \in \mathbb{N}\}$ denote the queue-length process embedded right after the departure of the n th customer, see [6]. Note that X_n has state-space $\{0, \dots, N-1\}$ as after the departure of a customer the system cannot be full.

Then $\{X_n : n \in \mathbb{N}\}$ is a Markov chain with transition matrix

$$P = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & \cdots & a_{N-2} & 1 - \sum_{k=0}^{N-2} a_k \\ a_0 & a_1 & a_2 & a_3 & \cdots & a_{N-2} & 1 - \sum_{k=0}^{N-2} a_k \\ 0 & a_0 & a_1 & a_2 & \cdots & a_{N-3} & 1 - \sum_{k=0}^{N-3} a_k \\ 0 & 0 & a_0 & a_1 & \cdots & a_{N-4} & 1 - \sum_{k=0}^{N-4} a_k \\ & & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & a_0 & 1 - a_0 \end{pmatrix}, \quad (4)$$

where

$$a_k = \int_0^\infty \frac{(\lambda t)^k}{k!} e^{-\lambda t} dB(t), \quad k = 0, \dots, N-2. \quad (5)$$

In words, a_k is the probability of k Poisson arrivals during an $B(\cdot)$ distributed service time.

3 The Taylor Series Expansion Approach

In this section, we present the Taylor series approximation for the M/G/1/N queue.

Let $B(\cdot)$ have density mapping $b(\cdot)$. Let $\Theta = (a, b) \subset \mathbb{R}$, for $0 < a < b < \infty$.

(A) For $0 \leq k \leq N-2$ it holds that a_k is n -times differentiable with respect to θ on Θ .

Under (A) it holds that the first n derivatives of P exists. Let $P^{(k)}$ denote the k th order derivative of P with respect to θ , then it holds that

$$P^{(k)}(i, j) = \frac{d^{(k)}}{d\theta^{(k)}} P(i, j), \quad 0 \leq i, j \leq N-1,$$

or, more specifically,

$$P^{(k)} = \begin{pmatrix} a_0(k) & a_1(k) & a_2(k) & a_3(k) & \cdots & a_{N-2}(k) & -\sum_{j=0}^{N-2} a_j(k) \\ a_0(k) & a_1(k) & a_2(k) & a_3(k) & \cdots & a_{N-2}(k) & -\sum_{j=0}^{N-2} a_j(k) \\ 0 & a_0(k) & a_1(k) & a_2(k) & \cdots & a_{N-3}(k) & -\sum_{j=0}^{N-3} a_j(k) \\ 0 & 0 & a_0(k) & a_1(k) & \cdots & a_{N-4}(k) & -\sum_{j=0}^{N-4} a_j(k) \\ & & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & a_0(k) & -a_0(k) \end{pmatrix} \quad (6)$$

where

$$a_j(k) = \frac{d^k}{d\theta^k} a_j, \quad 0 \leq j \leq N-2.$$

Example 1 Consider the $M/D/1$ queue with arrival rate λ and deterministic service rate c . Then, a_k is given by the probability to see k arrivals in an time interval of length c :

$$a_k = \frac{(\lambda c)^k}{k!} e^{-\lambda c}.$$

Example 2 Consider the $M/G/1$ queue with arrival rate λ and Weibull- (μ, γ) -distributed service times, where the density function is given by

$$b(t) = \mu \gamma (\mu t)^{\gamma-1} e^{-(\mu t)^\gamma}, \quad t \geq 0.$$

Note that the mean of the Weibull- (μ, γ) -distribution is $1/\mu$ and that γ defines the shape of the distribution. In particular, $\gamma = 1$ yields the exponential distribution with rate μ . The influence of γ on the shape of the service time distribution is illustrated in more detail in Section 4.2. Then,

$$a_k = \frac{\gamma \mu^\gamma \lambda^k}{k!} \int_0^\infty t^{k+\gamma-1} e^{-\lambda t - (\mu t)^\gamma} dt.$$

Let π_θ denote the stationary distribution of the embedded chain, where θ denotes the parameter of interest, and denote the deviation matrix by D_θ defined by

$$D_\theta = \sum_{n=0}^{\infty} (P_\theta^n - \Pi_\theta),$$

where Π_θ is a square matrix with rows equal to π_θ^\top , with x^\top denoting the transposed of vector x . As shown in [9], for any finite-state aperiodic Markov chain the deviation matrix exists.

Theorem 1 Let $\theta \in \Theta$ and let $\Theta_0 \subset \Theta$ a closed interval with θ be an interior point such that the queue is stable on Θ_0 . Provided that the entries of P are n -times differentiable with respect to θ , let

$$K_\theta(n) = \sum_{\substack{1 \leq m \leq n \\ 1 \leq l_k \leq n \\ l_1 + \dots + l_m = n}} \frac{n!}{l_1! \dots l_m!} \prod_{k=1}^m \left(P_\theta^{(l_k)} D_\theta \right).$$

Then it holds that

$$\pi_\theta^{(n)} = \pi_\theta K_\theta(n).$$

Proof: We prove the theorem by induction. For $n = 1$, we have to show that $\pi'_\theta = \pi_\theta P'_\theta D_\theta$. By simple algebra, it holds that for Δ such that $\theta + \Delta \in \Theta$ that

$$\pi_{\theta+\Delta} - \pi_\theta = \pi_{\theta+\Delta} (P_{\theta+\Delta} - P_\theta) D_\theta,$$

see, e.g., [9] for a proof, which yields

$$\frac{1}{\Delta}(\pi_{\theta+\Delta} - \pi_\theta) = \pi_\theta \frac{1}{\Delta}(P_{\theta+\Delta} - P_\theta)D_\theta + (\pi_{\theta+\Delta} - \pi_\theta) \frac{1}{\Delta}(P_{\theta+\Delta} - P_\theta)D_\theta. \quad (7)$$

Element-wise differentiability of P implies that

$$\lim_{\Delta \rightarrow 0} \pi_\theta \frac{1}{\Delta}(P_{\theta+\Delta} - P_\theta)D_\theta = \pi_\theta P'_\theta D_\theta. \quad (8)$$

Since, Θ_0 is a compact neighborhood of θ , and π_θ is finite for any $\theta \in \Theta_0$, it holds that

$$\sup_{\theta \in \Theta_0} |\pi_\theta|$$

is finite. Moreover,

$$0 \leq |(\pi_{\theta+\Delta} - \pi_\theta)| \left| \frac{1}{\Delta}(P_{\theta+\Delta} - P_\theta)D_\theta \right| \leq \sup_{\theta \in \Theta_0} |\pi_\theta| \left| \frac{1}{\Delta}(P_{\theta+\Delta} - P_\theta)D_\theta \right|.$$

Element-wise differentiability of P then yields

$$\lim_{\Delta \rightarrow 0} \left| \frac{1}{\Delta}(P_{\theta+\Delta} - P_\theta)D_\theta \right| = 0,$$

which implies that the term on the righthand side of (7) tends to zero as Δ tends to zero. Hence, taking the limit for Δ to zero in (7) reduces to (8), which proves the claim for $n = 1$.

The proof for the general case follows by induction with respect to n like in conventional analysis. \square

Remark 1 *The result put forward in Theorem 1 appears to be a special case of Theorem 4 in [7]. However, the latter result was established for Markov chains with general state-space and in the general case the analysis of the differentiability of π_θ requires elaborate conditions concerning the geometric ergodicity of the Markov chain with respect to a particular type of norm corresponding to the Lyapunov function of the system. This results in conditions that are often hard to check, even for simple systems. For this reason, we provide in Theorem 1 a new, simple and self-contained proof that is tailored to the class of problems studied in this paper.*

The derivatives in Theorem 1 enjoy a recursive structure in the sense that a $(k + 1)$ -st order derivative is mainly constituted out of information already provided by the k -th order derivative. The following lemma provides the exact statement.

Lemma 1 *Under the conditions put forward in Theorem 1 it holds for $k < n$ that*

$$\pi_\theta^{(k+1)} = \sum_{m=0}^k \binom{k+1}{m} \pi_\theta^{(m)} P_\theta^{(k+1-m)} D_\theta.$$

Example 3 For ease of reference we will provide in the following an explicit representation of the first derivatives of π_θ :

$$\pi_\theta^{(1)} = \pi_\theta P_\theta^{(1)} D_\theta$$

and

$$\pi_\theta^{(2)} = \pi_\theta P_\theta^{(2)} D_\theta + 2\pi_\theta (P_\theta^{(1)} D_\theta)^2.$$

Elaborating on the recursive formula for higher order derivatives in Lemma 1, the second order derivative can be written as

$$\pi_\theta^{(2)} = \pi_\theta P_\theta^{(2)} D_\theta + 2\pi_\theta^{(1)} P_\theta^{(1)} D_\theta.$$

In the same vein, we obtain for the third order derivative

$$\pi_\theta^{(3)} = \pi_\theta P_\theta^{(3)} D_\theta + 3\pi_\theta^{(2)} P_\theta^{(1)} D_\theta + 3\pi_\theta^{(1)} P_\theta^{(2)} D_\theta,$$

and

$$\pi_\theta^{(4)} = \pi_\theta P_\theta^{(4)} D_\theta + 4\pi_\theta^{(3)} P_\theta^{(1)} D_\theta + 6\pi_\theta^{(2)} P_\theta^{(2)} D_\theta + 4\pi_\theta^{(1)} P_\theta^{(3)} D_\theta.$$

A Taylor polynomial yields an approximation and the error introduced by this approximation can be expressed by the Lagrange form of the remainder as follows

$$r_\theta(k, \Delta) = \int_0^\Delta \frac{x^k}{k!} \pi_{\theta+x}^{(k+1)} dx. \quad (9)$$

From a numerical point of view the above expression is rather pointless as, by Theorem 1, it holds that

$$\pi_{\theta+x}^{(k+1)} = \pi_{\theta+\Delta} K_{\theta+\Delta}(k+1)$$

which implies that for computing the remainder we already have to know the very entity we want to approximate, namely, π_η for $\eta \in [\theta, \theta + \eta]$. To overcome this drawback we will present an alternative form for the remainder term.

The basic idea is that analyticity of π_θ implies that of $\pi_\theta^{(k)}$ for all k and we can again use a Taylor series to approximate $\pi_{\theta+x}^{(k+1)}$ in (9). By doing so we initiate the Taylor series in the tail of original Taylor series, and we expect that the error of this second Taylor approximation step is negligibly small. We explain this approach in the following.

Let

$$G_\theta(k, m, \delta) = \sum_{n=0}^m \frac{\delta^n}{n!} \pi_\theta^{(k+1+n)}$$

denote the Taylor polynomial of order m for $\pi_\theta^{(k+1)}$, i.e.,

$$\pi_{\theta+\delta}^{(k+1)} \approx G_\theta(k, m, \delta)$$

for m sufficiently large. Inserting the above approximation into (9) yields

$$\begin{aligned}
r_\theta(k, \Delta) &\approx \int_0^\Delta \frac{x^k}{k!} G_\theta(k, m, x) dx \\
&= \int_0^\Delta \frac{x^k}{k!} \sum_{n=0}^m \frac{x^n}{n!} \pi_\theta^{(k+1+n)} dx \\
&= \sum_{n=0}^m \frac{1}{k! n!} \pi_\theta^{(k+1+n)} \int_0^\Delta x^{k+n} dx \\
&= \sum_{n=0}^m \frac{\Delta^{k+n+1}}{k! n! (k+n+1)} \pi_\theta^{(k+1+n)}. \tag{10}
\end{aligned}$$

We denote by

$$g_\theta(k, m, \Delta) = \sum_{n=0}^m \frac{\Delta^{k+n+1}}{k! n! (k+n+1)} \pi_\theta^{(k+1+n)}$$

the expression for the approximation of the remainder term obtained from (10). Provided that $|g_\theta(k, m, \Delta) - r_\theta(k, \Delta)|$ is small for m small, we will use $g_\theta(k, m, \Delta)$ in our Taylor series approach to determine the order of the polynomial that is sufficient for achieving the desired precision of the approximation. As the following theorem shows, $g_\theta(k, m, \Delta)$ is of order $c\Delta^m r_\theta(k, \Delta)$ for some small constant c . In other words, letting, for example, $\Delta = 0.1$ and choosing $m = 3$, the error introduced by our approximation of the remainder term at k is typically smaller than $10^{-(5+k)}$.

In order to state the precise statement, we introduce the norm

$$\|x\| = \sum_{i=1}^n |x_i|$$

on \mathbb{R}^n .

Theorem 2 *Let $\theta \in \Theta$ be an interior point of Θ and let $\Delta > 0$ be such that $\theta + \Delta \in \Theta$. Assume that the entries of P are $(k+m+2)$ -times continuously differentiable with respect to θ on Θ . Suppose that a finite constant d exists such that*

$$d \geq \sup_{x \in [\theta, \theta + \Delta]} \|\pi_x^{(k+2+m)}\|,$$

then

$$|g_\theta(k, m, \Delta) - r_\theta(k, \Delta)| \leq d \frac{\Delta^{(m+k+2)}}{m! k! (m+k+2)}$$

Proof: Note that the Lagrange form of the remainder for $G_\theta(k, m, \delta)$ reads

$$\int_0^\delta \frac{u^m}{m!} \pi_{\theta+u}^{(k+m+2)} du.$$

Applying the norm $\|\cdot\|$ and using the bound d , yields

$$\begin{aligned} |g_\theta(k, m, \Delta) - r_\theta(k, \Delta)| &\leq \left| \int_0^\Delta \frac{x^k}{k!} \left(\int_0^x \frac{u^m}{m!} \pi_{\theta+u}^{(k+m+2)} du \right) dx \right| \\ &\leq d \int_0^\Delta \frac{x^k}{k!} \left(\int_0^x \frac{u^m}{m!} du \right) dx \\ &= d \frac{\Delta^{m+k+2}}{m! k! (m+k+2)}, \end{aligned}$$

which proves the claim. \square

In the numerical examples presented in the following sections, we will show that choosing $m = 2$ already yields a sufficient precision for approximating the remainder term.

Remark 2 *Taylor series like approaches for performance approximation have been studied in the literature before, see, e.g. [3, 4]. However, no a priori knowledge on the quality of the approximation of these approach could be established.*

Remark 3 *The Taylor series approximation developed above applies to differentiable Markov kernels. This extends the case of linear θ dependence that has been studied in the literature so far; see, for example, [1, 14, 21]. More specifically, for linear perturbations the standard assumption is that P_θ , the Markov kernel of the embedded jump chain, is of form*

$$P_\theta = \theta P + (1 - \theta) \hat{P}, \quad \theta \in [0, 1]. \quad (11)$$

In this case, P_θ has derivative $P'_\theta = P - \hat{P}$ and elements of Taylor series are of simple algebraical form

$$\pi_\theta^{(n)} = \pi_\theta((P - \hat{P})D_\theta)^n, \quad (12)$$

where D_θ is the deviation matrix associated with P_θ . See [1] for the denumerable state-space case and [8] for the general state-space case. Moreover, it has been shown in [9] that the remainder term can be bounded in terms of π_θ , P , \hat{P} and D_θ . This algorithm has earned its merits as the bound of the remainder term is numerically efficient and due to its simple algebraical form, the Taylor series can be easily computed. Extensions of this basic algorithm to continuous time processes are provided in [7, 15].

In contrast to the model in (11), the Taylor series expansions established in this section applies to non-linear perturbations. The case of a non-linear perturbation hasn't been studied in the literature so far. This case is significantly more difficult than the linear case as in this case all higher-order derivatives of P_θ maybe different from zero. Moreover, the elements of the Taylor series in the linear case are of rather simple form, compare the expression for $\pi_\theta^{(n)}$ in Theorem 1 with (12). This stems from the fact that in the linear case all but the first derivative of P with respect to θ are zero.

4 Applications to the M/G/1/N Queue

In this section we present the two numerical examples. In the first one, we let the distribution of the service times be deterministic, with θ denoting the deterministic service time, and in the second one we assume Weibull distributed service times, where θ is the shape parameter of the distribution. As performance measure we focus on the blocking probability $\pi_\theta^*(N)$, which is due the fact that customers arrive according to a Poisson arrival stream, equal to the probability that an arriving customer is lost due to no available free waiting space. Let $E[B(\theta)]$ denote the mean service time depending on θ and assume that the arrival rate λ is independent of θ . Then, the traffic rate is given by

$$\rho(\theta) = \lambda E[B(\theta)].$$

Recall, that by (2), it holds that

$$\pi_{\theta+\Delta}^*(N) = \frac{\pi_{\theta+\Delta}(0) + \rho(\theta + \Delta) - 1}{\pi_{\theta+\Delta}(0) + \rho(\theta + \Delta)}. \quad (13)$$

Inserting our Taylor series expansion for $\pi_{\theta+\Delta}(0)$ into the above expression yields a functional representation of $\pi_{\theta+\Delta}^*(N)$ as function in Δ . Elaborating on (2), a similar procedure leads to a functional representation of the mean queue length and via Little's law to one of the stationary waiting time.

4.1 Parameter Sensitivity: The M/D/1/N Queue

Consider the M/D/1 queue with arrival rate λ and deterministic service time $c = \theta$. The elements of P are provided in Example 4.1.

Lemma 2 *For the M/D/1/N queue, P is infinitely often differentiable with respect to c .*

Proof: By (6) differentiability properties of P can be deduced from that of the α_j entries. By Example 1, all higher-order derivatives exist for a_j , which proves the claim. \square

The following numerical examples illustrate our approach. In all numerical examples we have set $\lambda = 1$ and $N = 5$. We first illustrate the numerical behavior of $\pi_\theta^{(n)}(i)$ for $0 \leq i \leq N - 1$ and $n = 1, 2, 3$, for the stationary distribution of the embedded jump chain of the M/D/1/N queue.

As a first example, we apply the Taylor series of order 1, i.e.,

$$\begin{aligned} H_\theta(1, \Delta) &= \pi_\theta + \Delta \pi'_\theta \\ &= \pi_\theta + \Delta \pi_\theta P_\theta^{(1)} D_\theta, \end{aligned}$$

see Example 3. and we plot in Figure 1 the relative error given by

$$\frac{H_\theta(1, \Delta)(i) - \pi_{\theta+\Delta}(i)}{\pi_{\theta+\Delta}(i)}$$

for each element $i = 0, \dots, 4$. In words, the relative error times the factor 100 yields the error percentage of the Taylor series approximation. For Figure 1, we let $\theta = 1$ and vary Δ by at most 10 percent of θ , i.e., $0 \leq \Delta \leq \delta = 0.1$. Note that in this situation the traffic load of the system is one but, due to the fact that we consider a loss system, stability is still guaranteed.




Figure 1: The relative error in predicting $\pi_{1+\Delta}$ by $H_1(1, \Delta)$ for $\rho = 1$.

We now repeat this example, but we apply a Taylor series of order 2, i.e.,

$$\begin{aligned} H_\theta(2, \Delta) &= \pi_\theta + \Delta \pi_\theta^{(1)} + \frac{\Delta^2}{2} \pi_\theta^{(2)} \\ &= \pi_\theta + \Delta \pi_\theta P_\theta^{(1)} D_\theta + \frac{\Delta^2}{2} \pi_\theta P_\theta^{(2)} D_\theta + \Delta^2 \pi_\theta ((P_\theta^{(1)} D_\theta)^{(2)}), \end{aligned}$$

see Example 3, and we plot in Figure 2 the relative error given by

$$\frac{H_\theta(2, \Delta)(i) - \pi_{\theta+\Delta}(i)}{\pi_{\theta+\Delta}(i)}$$

for each element $i = 0, \dots, 4$. As for Figure 1, we let $\theta = 1$ and vary Δ by at most 10 percent of θ , i.e., $0 \leq \Delta \leq 0.1$.




Figure 2: The relative error in predicting $\pi_{1+\Delta}$ by $H_1(2, \Delta)$ for $\rho = 1$.

We conclude this series of examples by plotting the relative error of the Taylor series expansion for order 3 in Figure 3, where

$$\begin{aligned}
 H_\theta(3, \Delta) &= \pi_\theta + \Delta \pi_\theta^{(1)} + \frac{\Delta^2}{2} \pi_\theta^{(2)} + \frac{\Delta^3}{6} \pi_\theta^{(3)} \\
 &= \pi_\theta + \Delta \pi_\theta P_\theta^{(1)} D_\theta + \frac{\Delta^2}{2} \pi_\theta P_\theta^{(2)} D_\theta + \Delta^2 \pi_\theta (P_\theta^{(1)} D_\theta)^2 \\
 &\quad + \frac{\Delta^3}{6} \left(\pi_\theta P_\theta^{(3)} D_\theta + 3\pi_\theta^{(2)} P_\theta^{(1)} D_\theta + 3\pi_\theta^{(1)} P_\theta^{(2)} D_\theta \right),
 \end{aligned}$$

see Example 3.

Comparing Figure 1 to Figure 3 one notes that the state i which yields the dominant relative error changes with the order of the derivative. More specifically, while for the first order derivative, $\pi'_\theta(0)$ yields the dominant error in Figure 1, the dominant error for the second order derivative stems from state $i = 4$, see Figure 2. As can be seen from the figures, a third order Taylor series yields a remarkably good approximation of the stationary probabilities through the range $\theta \pm 0.1$. It is also worth noting that the quality of approximation increases with increasing traffic load. This is of particular interest as standard numerical methods for approximation the M/G/1/N queue are restricted to the case of $\rho < 1$.




Figure 3: The relative error in predicting $\pi_{1+\Delta}$ by $H_1(3, \Delta)$ for $\rho = 1$.

We now turn to the blocking probability. Starting point is the expression for the loss probability in (13). The traffic rate is given by $\rho(\theta + \Delta) = \lambda(\theta + \Delta)$ and $\pi_{\theta+\Delta}(0)$ is approximated via a Taylor series polynomial of degree k , i.e., we replace $\pi_{\theta+\Delta}(0)$ by $H_\theta(k, \Delta)(0)$. Figure 4 shows the relative absolute error in medium traffic, i.e., $\theta = 0.5$, for $k = 2, 3$. The approximation in case of saturation, i.e., $\rho(\theta) = 1$, is illustrated in Figure 5. Finally, in Figure 6 we show the behavior of our approximation for the case of over saturation, i.e., $\rho = 1.2$.

As can be seen from the figures, the approximation yields a satisfying precision in predicting the loss probability $\pi_{\theta+\Delta}^*(N)$ as a mapping of Δ in a range of Δ being 10 percent of θ .

We conclude the discussion of the M/D/1/N queue by providing a bound on the error of the Taylor series approximation for $\pi_{\theta+\Delta}^*(N)$. Suppose that for given order k of the Taylor approximation, the error is bounded by R for any Δ such that $|\Delta| \leq \delta$, i.e., assume that

$$|\pi_{\theta+\Delta}(0) - H_\theta(k, \Delta)| \leq R,$$

see Theorem 2. Replacing $\pi_{\theta+\Delta}(0)$ in (2) or (13) by $H_\theta(k, \Delta)$ and noting that $\rho(\theta + \Delta) = \lambda\theta + \Delta\lambda$, implies thus that the true value for $\pi_{\theta+\Delta}^*(N)$ is bounded by

$$\frac{H_\theta(k, \Delta) - R + \lambda\theta + \Delta\lambda - 1}{H_\theta(k, \Delta) - R + \lambda\theta + \Delta\lambda} \leq \pi_{\theta+\Delta}^*(N) \leq \frac{H_\theta(k, \Delta) + R + \lambda\theta + \Delta\lambda - 1}{H_\theta(k, \Delta) + R + \lambda\theta + \Delta\lambda}$$




Figure 4: The relative error in predicting the loss probability in medium traffic ($\rho = 0.5, \theta = 2$).

The numerical error can thus be bounded by

$$\begin{aligned} & \left| \frac{H_\theta(k, \Delta) + R + \lambda\theta + \Delta\lambda - 1}{H_\theta(k, \Delta) + R + \lambda\theta + \Delta\lambda} - \frac{H_\theta(k, \Delta) - R + \lambda\theta + \Delta\lambda - 1}{H_\theta(k, \Delta) - R + \lambda\theta + \Delta\lambda} \right| \\ &= \frac{2R}{(H_\theta(k, \Delta) + R + \lambda\theta + \Delta\lambda)(H_\theta(k, \Delta) - R + \lambda\theta + \Delta\lambda)} \end{aligned}$$

noting that $H_\theta(k, \Delta) \geq 0$ for all Δ , yields

$$\leq \frac{2R}{(R + \lambda\theta + \Delta\lambda)(-R + \lambda\theta + \Delta\lambda)} = \frac{2R}{\lambda^2(\theta - \Delta)^2 - R^2}$$

and from $|\Delta| \leq \delta$ it follows

$$\leq \frac{2R}{\lambda^2(\theta - \delta)^2 - R^2}.$$

We summarize our analysis in the following lemma.

Lemma 3 *Consider the $M/D/1/N$ queue with arrival rate λ and deterministic service time θ . Suppose that for k it holds for $|\Delta| \leq \delta$ that*

$$|\pi_{\theta+\Delta}(0) - H_\theta(k, \Delta)| \leq R_\theta(k, \delta),$$




Figure 5: The relative error in predicting the loss probability in saturated traffic ($\rho = 1, \theta = 1$).

then

$$\sup_{|\Delta| \leq \delta} \left| \pi_{\theta+\Delta}^*(N) - \frac{H_\theta(k, \Delta) + \lambda(\theta + \Delta) - 1}{H_\theta(k, \Delta) + \lambda(\theta + \Delta)} \right| \leq \frac{2R_\theta(k, \delta)}{\lambda^2(\theta - \delta)^2 - (R_\theta(k, \delta))^2}.$$

We conclude this section with a discussion of the numerical bound on the error provided in Lemma 3 and Theorem 2. To this end we consider the Taylor series approximation of degree $k = 2$ for the blocking probability. Figure 7 plots the true remainder term against the approximation of the remainder term obtained from inserting the expression in Theorem 2 into the bound provided in Lemma 3, where we have chosen $m = 2$. As can be seen from Figure 7, the approximation of the remainder term yields good results for small values of Δ . For example, the approximative remainder term indicates that a Taylor series of degree 2 yields a maximal error of 5×10^{-4} in predicting blocking probability, whereas the true error is no greater than 3×10^{-4} .

4.2 Distributional Sensitivity: The M/Weibull/1/N Queue

In this section we illustrate the use of the Taylor series for the M/G/1/N queue where the service time distribution is given by the Weibull distribution, see Example 2. We let $\theta = \gamma$, which yields that for $\theta = 1$ the service distribution




Figure 6: The relative error in predicting the loss probability in over-saturated traffic ($\rho = 1.2, \theta = 0.833$).

$B(t)$ yields the exponential distribution with mean $1/\mu$. Note that for $\theta > 1$, $B(t)$ is a heavy tailed distribution, whereas for $\theta < 1$ it becomes light tailed. Varying θ around one, provides a sensitivity analysis with respect to the shape of the distribution.

Lemma 4 *For the M/Weibull/1/N queue, P is infinitely often differentiable with respect to γ .*

Proof: By (6) differentiability properties of P can be deduced from that of the α_j entries. By Example 2, all higher-order derivatives exist for a_j , which proves the claim. \square

In the following we will let $\lambda = \mu = 1$, which implies $\rho = 1$ and independent of θ . We let $|\Delta| \leq \delta = 0.1$ and we illustrate the effect a change of θ by $\pm\Delta$ has on the density in Figure 8.

Like for the M/D/1/N case, we present in the following the relative errors for predicting $\pi_{\theta+\Delta}^*$ for various order of the Taylor polynomial and traffic rate one, where we restrict Δ to positive values, i.e., $0 \leq \Delta \leq \delta$. Figure 9 illustrates the relative error of the Taylor series approximation of degree 1. The relative error of the Taylor series approximation of degree 2 is shown in Figure 10. Eventually, we show relative error of the Taylor series approximation of degree 3 is shown in Figure 11.




Figure 7: The remainder term vs. the bound for the remainder at $k = 2$ and $m = 2$.

Comparing the numerical results with those for the M/D/1/N case, one notes that the qualitative behavior of the relative error is similar but that the effect of perturbing the value of the mean service time in the M/G/1/N queue is better predictable than that of perturbing the shape of the service time distribution.

We conclude the example with presenting the absolute relative error for predicting the blocking probability in the M/G/1/N queue for various traffic rates, where for $\rho = 1.2$ we plot only the range $|\Delta| \leq 0.05$ for better visibility of the curve.

We now turn to the bound on the remainder term. Since for the M/W/1/N model the traffic rate $\rho = \lambda/\mu$ is independent of θ , we obtain the following adaptation of Lemma 3 to the M/W/1/N queue.

Lemma 5 *Consider the M/W/1/N queue with arrival rate λ , service rate μ , and shape parameter θ . Denote the traffic rate by $\rho = \lambda/\mu$. Suppose that for k it holds for $|\Delta| \leq \delta$ that*

$$|\pi_{\theta+\Delta}(0) - H_{\theta}(k, \Delta)| \leq R_{\theta}(k, \delta),$$

then

$$\sup_{|\Delta| \leq \delta} \left| \pi_{\theta+\Delta}^*(N) - \frac{H_{\theta}(k, \Delta) + \rho - 1}{H_{\theta}(k, \Delta) + \rho} \right| \leq \frac{2 R_{\theta}(k, \delta)}{\rho^2 - (R_{\theta}(k, \delta))^2}.$$




Figure 8: The extreme values of the density

We conclude this section with a discussion of the numerical bound on the error provided in Lemma 3 and Theorem 2. To this end we consider the Taylor series approximation of degree $k = 2$ for the blocking probability. Figure 15 plots the true remainder term against the approximation of the remainder term obtained from inserting the expression in Theorem 2 into the bound provided in Lemma 5, where we have chosen $m = 2$. As can be seen from Figure 15, the approximation of the remainder term yields good results for small values of Δ . For example, the approximative remainder term indicates that a Taylor series of degree 2 yields a maximal error of 4.5×10^{-5} in predicting blocking probability, whereas the true error is no greater than 3.7×10^{-5} .

5 Conclusion

We have presented a new approach to the functional approximation of finite queues. As illustrated by the numerical examples for the M/G/1/N queue, the convergence rate of the Taylor series is such that already a Taylor polynomial of




Figure 9: The relative error in predicting $\pi_{1+\Delta}$ by $H_1(1, \Delta)$ for $\rho = 1$.

degree 2 or 3 yields good numerical results. We established an approximation for the remainder term of the Taylor series that provides an efficient way of computing (approximately) the remainder term and thereby provides an algorithmic way of deciding which order of the Taylor polynomial is sufficient to achieve a desired precision of the approximation. This implies the proposed Taylor series approximation can be of practical value. Future research will be on investigating the behavior of the series expansion for multi-server queues.

References

- [1] X.-R. Cao, The Maclaurin Series for performance functions of Markov chains. *Advances in Applied Probability* **30** (1998) 676–692.
- [2] M. Fu and J.-Q. Hu, *Conditional Monte Carlo*. Kluwer Academic, Boston, 1997.
- [3] M. Girish and J.Q. Hu, Higher order approximations for tandem queueing networks. *Queueing Systems* **22** (1996) 249–276.
- [4] M. Girish and J.Q. Hu, An interpolation approximation for the G/G/1 queue based on multipoint Padé approximation. *Queueing Systems* **26** (1997) 269–284.




Figure 10: The relative error in predicting $\pi_{1+\Delta}$ by $H_1(2, \Delta)$ for $\rho = 1$.

- [5] P. Glasserman, *Gradient Estimation via Perturbation Analysis*. Kluwer Academic Publishers, Boston, 1991.
- [6] D. Gross and C. Harris, *Fundamentals of Queueing Theory*, Wiley, 1985.
- [7] B. Heidergott and A. Hordijk, Taylor series expansions for stationary Markov chains. *Advances in Applied Probability* **35** (2003) 1046–1070.
- [8] B. Heidergott, A. Hordijk and N. Leder, Series expansions for continuous-time Markov chains. *Operations Research* **58** (2010) 756–767.
- [9] B. Heidergott, A. Hordijk and M. van Uitert, Series expansions for finite-state Markov chains. *Probability in Engineering and Informational Sciences* **21** (2007) 381–400.
- [10] Y.-C. Ho and X.-R. Cao, *Perturbation Analysis of Discrete Event Dynamic Systems*. Kluwer Academic, Boston, 1991.
- [11] A. Hordijk and F. M. Spieksma, *A New Formula for the Deviation Matrix*, Chapter 36 in *Probability, Statistics and Optimization* (F.P. Kelly, ed.) Wiley, 1994.
- [12] N.V. Kartashov, *Strong Stable Markov Chains*, Edition VSP, Utrecht TBIMC Scientific Publishers, Kiev, 1996.




Figure 11: The relative error in predicting $\pi_{1+\Delta}$ by $H_1(3, \Delta)$ for $\rho = 1$.

- [13] D. Kendall, Stochastic processes occurring in the theory of queues and their analysis by the method of embedded Markov chains. *Ann. Math. Statist.* **24** (1953) 338-354.
- [14] S. Kirkland, M. Neumann and B. Shader, Application of Paz's inequality to perturbation bounds for Markov chains. *Linear Algebra and its Applications* (1998) **268** 183-196.
- [15] N. Leder, B. Heidergott and A. Hordijk, An approximation approach for the deviation matrix of continuous-time Markov processes with applications to Markov decision theory. *Operations Research* **58** (2010) 918-932.
- [16] Z. Mouhoubi and D. Aïssani, New perturbation bounds for denumerable Markov chains. *Linear Algebra and its Applications* **432** (2010) 1627-1649.
- [17] G. Pflug, *Optimisation of Stochastic Models*. Kluwer Academic, Boston, 1996.
- [18] M. Reiman and A. Weiss, Sensitivity Analysis for Simulations via Likelihood Ratios. *Operations Research*, **37** (1989) 830-844.
- [19] B. Ripley, *Stochastic Simulation*. Wiley & Sons, New York, 1987.




Figure 12: The relative error in predicting the loss probability in dense traffic ($\rho = 0.7$)

- [20] R. Rubinstein and A. Shapiro, *Discrete Event Systems: Sensitivity Analysis and Optimization by the Score Function Method*. Wiley, 1993.
- [21] E. Schweitzer, Perturbation theory and finite Markov chains. *Journal of Applied Probability* **5** (1968) 401-413.
- [22] J. M. Smith, Optimal Design and Performance Modelling of $M/G/1/K$ Queueing Systems. *Mathematical and Computer Modelling* **39** (2004) 1049–1081.
- [23] H. Tijms, Heuristics for finite-buffer queues. *Probability in the Engineering and Informational Sciences* **6** (1992) 277–285.
- [24] H. Tijms, *Stochastic Modeling and Analysis*, Wiley, New York, 1986.




Figure 13: The relative error in predicting the loss probability in saturated traffic ($\rho = 1$)




Figure 14: The relative error in predicting the loss probability in over saturated traffic ($\rho = 1.2$)




Figure 15: The remainder term vs. the bound for the remainder at $k = 2$ and $m = 2$.

2007-1	M. Francesca Cracolici Miranda Cuffaro Peter Nijkamp	Geographical distribution of enemployment: An analysis of provincial differences in Italy, 21 p.
2007-2	Daniel Leliefeld Evgenia Motchenkova	To protect in order to serve, adverse effects of leniency programs in view of industry asymmetry, 29 p.
2007-3	M.C. Wassenaar E. Dijkgraaf R.H.J.M. Gradus	Contracting out: Dutch municipalities reject the solution for the VAT-distortion, 24 p.
2007-4	R.S. Halbersma M.C. Mikkers E. Motchenkova I. Seinen	Market structure and hospital-insurer bargaining in the Netherlands, 20 p.
2007-5	Bas P. Singer Bart A.G. Bossink Herman J.M. Vande Putte	Corporate Real estate and competitive strategy, 27 p.
2007-6	Dorien Kooij Annet de Lange Paul Jansen Josje Dijkers	Older workers' motivation to continue to work: Five meanings of age. A conceptual review, 46 p.
2007-7	Stella Flytzani Peter Nijkamp	Locus of control and cross-cultural adjustment of expatriate managers, 16 p.
2007-8	Tibert Verhagen Willemijn van Dolen	Explaining online purchase intentions: A multi-channel store image perspective, 28 p.
2007-9	Patrizia Riganti Peter Nijkamp	Congestion in popular tourist areas: A multi-attribute experimental choice analysis of willingness-to-wait in Amsterdam, 21 p.
2007-10	Tüzin Baycan-Levent Peter Nijkamp	Critical success factors in planning and management of urban green spaces in Europe, 14 p.
2007-11	Tüzin Baycan-Levent Peter Nijkamp	Migrant entrepreneurship in a diverse Europe: In search of sustainable development, 18 p.
2007-12	Tüzin Baycan-Levent Peter Nijkamp Mediha Sahin	New orientations in ethnic entrepreneurship: Motivation, goals and strategies in new generation ethnic entrepreneurs, 22 p.
2007-13	Miranda Cuffaro Maria Francesca Cracolici Peter Nijkamp	Measuring the performance of Italian regions on social and economic dimensions, 20 p.

2007-14	Tüzin Baycan-Levent Peter Nijkamp	Characteristics of migrant entrepreneurship in Europe, 14 p.
2007-15	Maria Teresa Borzacchiello Peter Nijkamp Eric Koomen	Accessibility and urban development: A grid-based comparative statistical analysis of Dutch cities, 22 p.
2007-16	Tibert Verhagen Selmar Meents	A framework for developing semantic differentials in IS research: Assessing the meaning of electronic marketplace quality (EMQ), 64 p.
2007-17	Aliye Ahu Gülümser Tüzin Baycan Levent Peter Nijkamp	Changing trends in rural self-employment in Europe, 34 p.
2007-18	Laura de Dominicis Raymond J.G.M. Florax Henri L.F. de Groot	De ruimtelijke verdeling van economische activiteit: Agglomeratie- en locatiepatronen in Nederland, 35 p.
2007-19	E. Dijkgraaf R.H.J.M. Gradus	How to get increasing competition in the Dutch refuse collection market? 15 p.

2008-1	Maria T. Borzacchiello Irene Casas Biagio Ciuffo Peter Nijkamp	Geo-ICT in Transportation Science, 25 p.
2008-2	Maura Soekijad Jeroen Walschots Marleen Huysman	Congestion at the floating road? Negotiation in networked innovation, 38 p.
2008-3	Marlous Agterberg Bart van den Hooff Marleen Huysman Maura Soekijad	Keeping the wheels turning: Multi-level dynamics in organizing networks of practice, 47 p.
2008-4	Marlous Agterberg Marleen Huysman Bart van den Hooff	Leadership in online knowledge networks: Challenges and coping strategies in a network of practice, 36 p.
2008-5	Bernd Heidergott Haralambie Leahu	Differentiability of product measures, 35 p.
2008-6	Tibert Verhagen Frans Feldberg Bart van den Hooff Selmar Meents	Explaining user adoption of virtual worlds: towards a multipurpose motivational model, 37 p.
2008-7	Masagus M. Ridhwan Peter Nijkamp Piet Rietveld Henri L.F. de Groot	Regional development and monetary policy. A review of the role of monetary unions, capital mobility and locational effects, 27 p.
2008-8	Selmar Meents Tibert Verhagen	Investigating the impact of C2C electronic marketplace quality on trust, 69 p.
2008-9	Junbo Yu Peter Nijkamp	China's prospects as an innovative country: An industrial economics perspective, 27 p
2008-10	Junbo Yu Peter Nijkamp	Ownership, r&d and productivity change: Assessing the catch-up in China's high-tech industries, 31 p
2008-11	Elbert Dijkgraaf Raymond Gradus	Environmental activism and dynamics of unit-based pricing systems, 18 p.
2008-12	Mark J. Koetse Jan Rouwendal	Transport and welfare consequences of infrastructure investment: A case study for the Betuweroute, 24 p
2008-13	Marc D. Bahlmann Marleen H. Huysman Tom Elfring Peter Groenewegen	Clusters as vehicles for entrepreneurial innovation and new idea generation – a critical assessment
2008-14	Soushi Suzuki Peter Nijkamp	A generalized goals-achievement model in data envelopment analysis: An application to efficiency improvement in local government finance in Japan, 24 p.
2008-15	Tüzin Baycan-Levent	External orientation of second generation migrant entrepreneurs. A sectoral

	Peter Nijkamp Mediha Sahin	study on Amsterdam, 33 p.
2008-16	Enno Masurel	Local shopkeepers' associations and ethnic minority entrepreneurs, 21 p.
2008-17	Frank Frößler Boriana Rukanova Stefan Klein Allen Higgins Yao-Hua Tan	Inter-organisational network formation and sense-making: Initiation and management of a living lab, 25 p.
2008-18	Peter Nijkamp Frank Zwetsloot Sander van der Wal	A meta-multicriteria analysis of innovation and growth potentials of European regions, 20 p.
2008-19	Junbo Yu Roger R. Stough Peter Nijkamp	Governing technological entrepreneurship in China and the West, 21 p.
2008-20	Maria T. Borzacchiello Peter Nijkamp Henk J. Scholten	A logistic regression model for explaining urban development on the basis of accessibility: a case study of Naples, 13 p.
2008-21	Marius Ooms	Trends in applied econometrics software development 1985-2008, an analysis of Journal of Applied Econometrics research articles, software reviews, data and code, 30 p.
2008-22	Aliye Ahu Gülümser Tüzin Baycan-Levent Peter Nijkamp	Changing trends in rural self-employment in Europe and Turkey, 20 p.
2008-23	Patricia van Hemert Peter Nijkamp	Thematic research prioritization in the EU and the Netherlands: an assessment on the basis of content analysis, 30 p.
2008-24	Jasper Dekkers Eric Koomen	Valuation of open space. Hedonic house price analysis in the Dutch Randstad region, 19 p.

2009-1	Boriana Rukanova Rolf T. Wignand Yao-Hua Tan	From national to supranational government inter-organizational systems: An extended typology, 33 p.
2009-2	Marc D. Bahlmann Marleen H. Huysman Tom Elfring Peter Groenewegen	Global Pipelines or global buzz? A micro-level approach towards the knowledge-based view of clusters, 33 p.
2009-3	Julie E. Ferguson Marleen H. Huysman	Between ambition and approach: Towards sustainable knowledge management in development organizations, 33 p.
2009-4	Mark G. Leijssen	Why empirical cost functions get scale economies wrong, 11 p.
2009-5	Peter Nijkamp Galit Cohen-Blankshtain	The importance of ICT for cities: e-governance and cyber perceptions, 14 p.
2009-6	Eric de Noronha Vaz Mário Caetano Peter Nijkamp	Trapped between antiquity and urbanism. A multi-criteria assessment model of the greater Cairo metropolitan area, 22 p.
2009-7	Eric de Noronha Vaz Teresa de Noronha Vaz Peter Nijkamp	Spatial analysis for policy evaluation of the rural world: Portuguese agriculture in the last decade, 16 p.
2009-8	Teresa de Noronha Vaz Peter Nijkamp	Multitasking in the rural world: Technological change and sustainability, 20 p.
2009-9	Maria Teresa Borzacchiello Vincenzo Torrieri Peter Nijkamp	An operational information systems architecture for assessing sustainable transportation planning: Principles and design, 17 p.
2009-10	Vincenzo Del Giudice Pierfrancesco De Paola Francesca Torrieri Francesca Pagliari Peter Nijkamp	A decision support system for real estate investment choice, 16 p.
2009-11	Miruna Mazurencu Marinescu Peter Nijkamp	IT companies in rough seas: Predictive factors for bankruptcy risk in Romania, 13 p.
2009-12	Boriana Rukanova Helle Zinner Hendriksen Eveline van Stijn Yao-Hua Tan	Bringing is innovation in a highly-regulated environment: A collective action perspective, 33 p.
2009-13	Patricia van Hemert Peter Nijkamp Jolanda Verbraak	Evaluating social science and humanities knowledge production: an exploratory analysis of dynamics in science systems, 20 p.

2009-14	Roberto Patuelli Aura Reggiani Peter Nijkamp Norbert Schanne	Neural networks for cross-sectional employment forecasts: A comparison of model specifications for Germany, 15 p.
2009-15	André de Waal Karima Kourtit Peter Nijkamp	The relationship between the level of completeness of a strategic performance management system and perceived advantages and disadvantages, 19 p.
2009-16	Vincenzo Punzo Vincenzo Torrieri Maria Teresa Borzacchiello Biagio Ciuffo Peter Nijkamp	Modelling intermodal re-balance and integration: planning a sub-lagoon tube for Venezia, 24 p.
2009-17	Peter Nijkamp Roger Stough Mediha Sahin	Impact of social and human capital on business performance of migrant entrepreneurs – a comparative Dutch-US study, 31 p.
2009-18	Dres Creal	A survey of sequential Monte Carlo methods for economics and finance, 54 p.
2009-19	Karima Kourtit André de Waal	Strategic performance management in practice: Advantages, disadvantages and reasons for use, 15 p.
2009-20	Karima Kourtit André de Waal Peter Nijkamp	Strategic performance management and creative industry, 17 p.
2009-21	Eric de Noronha Vaz Peter Nijkamp	Historico-cultural sustainability and urban dynamics – a geo-information science approach to the Algarve area, 25 p.
2009-22	Roberta Capello Peter Nijkamp	Regional growth and development theories revisited, 19 p.
2009-23	M. Francesca Cracolici Miranda Cuffaro Peter Nijkamp	Tourism sustainability and economic efficiency – a statistical analysis of Italian provinces, 14 p.
2009-24	Caroline A. Rodenburg Peter Nijkamp Henri L.F. de Groot Erik T. Verhoef	Valuation of multifunctional land use by commercial investors: A case study on the Amsterdam Zuidas mega-project, 21 p.
2009-25	Katrin Oltmer Peter Nijkamp Raymond Florax Floor Brouwer	Sustainability and agri-environmental policy in the European Union: A meta-analytic investigation, 26 p.
2009-26	Francesca Torrieri Peter Nijkamp	Scenario analysis in spatial impact assessment: A methodological approach, 20 p.
2009-27	Aliye Ahu Gülümser Tüzin Baycan-Levent Peter Nijkamp	Beauty is in the eyes of the beholder: A logistic regression analysis of sustainability and locality as competitive vehicles for human settlements, 14 p.

2009-28	Marco Percoco Peter Nijkamp	Individual time preferences and social discounting in environmental projects, 24 p.
2009-29	Peter Nijkamp Maria Abreu	Regional development theory, 12 p.
2009-30	Tüzin Baycan-Levent Peter Nijkamp	7 FAQs in urban planning, 22 p.
2009-31	Aliye Ahu Gülümser Tüzin Baycan-Levent Peter Nijkamp	Turkey's rurality: A comparative analysis at the EU level, 22 p.
2009-32	Frank Bruinsma Karima Kourtit Peter Nijkamp	An agent-based decision support model for the development of e-services in the tourist sector, 21 p.
2009-33	Mediha Sahin Peter Nijkamp Marius Rietdijk	Cultural diversity and urban innovativeness: Personal and business characteristics of urban migrant entrepreneurs, 27 p.
2009-34	Peter Nijkamp Mediha Sahin	Performance indicators of urban migrant entrepreneurship in the Netherlands, 28 p.
2009-35	Manfred M. Fischer Peter Nijkamp	Entrepreneurship and regional development, 23 p.
2009-36	Faroek Lazrak Peter Nijkamp Piet Rietveld Jan Rouwendal	Cultural heritage and creative cities: An economic evaluation perspective, 20 p.
2009-37	Enno Masurel Peter Nijkamp	Bridging the gap between institutions of higher education and small and medium-size enterprises, 32 p.
2009-38	Francesca Medda Peter Nijkamp Piet Rietveld	Dynamic effects of external and private transport costs on urban shape: A morphogenetic perspective, 17 p.
2009-39	Roberta Capello Peter Nijkamp	Urban economics at a cross-yard: Recent theoretical and methodological directions and future challenges, 16 p.
2009-40	Enno Masurel Peter Nijkamp	The low participation of urban migrant entrepreneurs: Reasons and perceptions of weak institutional embeddedness, 23 p.
2009-41	Patricia van Hemert Peter Nijkamp	Knowledge investments, business R&D and innovativeness of countries. A qualitative meta-analytic comparison, 25 p.
2009-42	Teresa de Noronha Vaz Peter Nijkamp	Knowledge and innovation: The strings between global and local dimensions of sustainable growth, 16 p.
2009-43	Chiara M. Traversi Peter Nijkamp	Managing environmental risk in agriculture: A systematic perspective on the potential of quantitative policy-oriented risk valuation, 19 p.
2009-44	Sander de Leeuw	Logistics aspects of emergency preparedness in flood disaster prevention, 24 p.

Iris F.A. Vis
Sebastiaan B. Jonkman

- | | | |
|---------|---|---|
| 2009-45 | Eveline S. van Leeuwen
Peter Nijkamp | Social accounting matrices. The development and application of SAMs at the local level, 26 p. |
| 2009-46 | Tibert Verhagen
Willemijn van Dolen | The influence of online store characteristics on consumer impulsive decision-making: A model and empirical application, 33 p. |
| 2009-47 | Eveline van Leeuwen
Peter Nijkamp | A micro-simulation model for e-services in cultural heritage tourism, 23 p. |
| 2009-48 | Andrea Caragliu
Chiara Del Bo
Peter Nijkamp | Smart cities in Europe, 15 p. |
| 2009-49 | Farook Lazrak
Peter Nijkamp
Piet Rietveld
Jan Rouwendal | Cultural heritage: Hedonic prices for non-market values, 11 p. |
| 2009-50 | Eric de Noronha Vaz
João Pedro Bernardes
Peter Nijkamp | Past landscapes for the reconstruction of Roman land use: Eco-history tourism in the Algarve, 23 p. |
| 2009-51 | Eveline van Leeuwen
Peter Nijkamp
Teresa de Noronha Vaz | The Multi-functional use of urban green space, 12 p. |
| 2009-52 | Peter Bakker
Carl Koopmans
Peter Nijkamp | Appraisal of integrated transport policies, 20 p. |
| 2009-53 | Luca De Angelis
Leonard J. Paas | The dynamics analysis and prediction of stock markets through the latent Markov model, 29 p. |
| 2009-54 | Jan Anne Annema
Carl Koopmans | Een lastige praktijk: Ervaringen met waarderen van omgevingskwaliteit in de kosten-batenanalyse, 17 p. |
| 2009-55 | Bas Straathof
Gert-Jan Linders | Europe's internal market at fifty: Over the hill? 39 p. |
| 2009-56 | Joaquim A.S. Gromicho
Jelke J. van Hoorn
Francisco Saldanha-da-Gama
Gerrit T. Timmer | Exponentially better than brute force: solving the job-shop scheduling problem optimally by dynamic programming, 14 p. |
| 2009-57 | Carmen Lee
Roman Kraeussl
Leo Paas | The effect of anticipated and experienced regret and pride on investors' future selling decisions, 31 p. |
| 2009-58 | René Sitters | Efficient algorithms for average completion time scheduling, 17 p. |

2009-59

Masood Gheasi
Peter Nijkamp
Piet Rietveld

Migration and tourist flows, 20 p.

2010-1	Roberto Patuelli Norbert Schanne Daniel A. Griffith Peter Nijkamp	Persistent disparities in regional unemployment: Application of a spatial filtering approach to local labour markets in Germany, 28 p.
2010-2	Thomas de Graaff Ghebre Debrezion Piet Rietveld	Schaalsprong Almere. Het effect van bereikbaarheidsverbeteringen op de huizenprijzen in Almere, 22 p.
2010-3	John Steenbruggen Maria Teresa Borzacchiello Peter Nijkamp Henk Scholten	Real-time data from mobile phone networks for urban incidence and traffic management – a review of application and opportunities, 23 p.
2010-4	Marc D. Bahlmann Tom Elfring Peter Groenewegen Marleen H. Huysman	Does distance matter? An ego-network approach towards the knowledge-based theory of clusters, 31 p.
2010-5	Jelke J. van Hoorn	A note on the worst case complexity for the capacitated vehicle routing problem, 3 p.
2010-6	Mark G. Lijesen	Empirical applications of spatial competition; an interpretative literature review, 16 p.
2010-7	Carmen Lee Roman Kraeussl Leo Paas	Personality and investment: Personality differences affect investors' adaptation to losses, 28 p.
2010-8	Nahom Ghebrihiwet Evgenia Motchenkova	Leniency programs in the presence of judicial errors, 21 p.
2010-9	Meindert J. Flikkema Ard-Pieter de Man Matthijs Wolters	New trademark registration as an indicator of innovation: results of an explorative study of Benelux trademark data, 53 p.
2010-10	Jani Merikivi Tibert Verhagen Frans Feldberg	Having belief(s) in social virtual worlds: A decomposed approach, 37 p.
2010-11	Umut Kilingç	Price-cost markups and productivity dynamics of entrant plants, 34 p.
2010-12	Umut Kilingç	Measuring competition in a frictional economy, 39 p.

2011-1	Yoshifumi Takahashi Peter Nijkamp	Multifunctional agricultural land use in sustainable world, 25 p.
2011-2	Paulo A.L.D. Nunes Peter Nijkamp	Biodiversity: Economic perspectives, 37 p.
2011-3	Eric de Noronha Vaz Doan Nainggolan Peter Nijkamp Marco Painho	A complex spatial systems analysis of tourism and urban sprawl in the Algarve, 23 p.
2011-4	Karima Kourtit Peter Nijkamp	Strangers on the move. Ethnic entrepreneurs as urban change actors, 34 p.
2011-5	Manie Geyer Helen C. Coetzee Danie Du Plessis Ronnie Donaldson Peter Nijkamp	Recent business transformation in intermediate-sized cities in South Africa, 30 p.
2011-6	Aki Kangasharju Christophe Tavera Peter Nijkamp	Regional growth and unemployment. The validity of Okun's law for the Finnish regions, 17 p.
2011-7	Amitrajeet A. Batabyal Peter Nijkamp	A Schumpeterian model of entrepreneurship, innovation, and regional economic growth, 30 p.
2011-8	Aliye Ahu Akgün Tüzin Baycan Levent Peter Nijkamp	The engine of sustainable rural development: Embeddedness of entrepreneurs in rural Turkey, 17 p.
2011-9	Aliye Ahu Akgün Eveline van Leeuwen Peter Nijkamp	A systemic perspective on multi-stakeholder sustainable development strategies, 26 p.
2011-10	Tibert Verhagen Jaap van Nes Frans Feldberg Willemijn van Dolen	Virtual customer service agents: Using social presence and personalization to shape online service encounters, 48 p.
2011-11	Henk J. Scholten Maarten van der Vlist	De inrichting van crisisbeheersing, de relatie tussen besluitvorming en informatievoorziening. Casus: Warroom project Netcentrisch werken bij Rijkswaterstaat, 23 p.
2011-12	Tüzin Baycan Peter Nijkamp	A socio-economic impact analysis of cultural diversity, 22 p.
2011-13	Aliye Ahu Akgün Tüzin Baycan Peter Nijkamp	Repositioning rural areas as promising future hot spots, 22 p.
2011-14	Selmar Meents Tibert Verhagen Paul Vlaar	How sellers can stimulate purchasing in electronic marketplaces: Using information as a risk reduction signal, 29 p.

2011-15	Aliye Ahu Gülümser Tüzin Baycan-Levent Peter Nijkamp	Measuring regional creative capacity: A literature review for rural-specific approaches, 22 p.
2011-16	Frank Bruinsma Karima Kourtit Peter Nijkamp	Tourism, culture and e-services: Evaluation of e-services packages, 30 p.
2011-17	Peter Nijkamp Frank Bruinsma Karima Kourtit Eveline van Leeuwen	Supply of and demand for e-services in the cultural sector: Combining top-down and bottom-up perspectives, 16 p.
2011-18	Eveline van Leeuwen Peter Nijkamp Piet Rietveld	Climate change: From global concern to regional challenge, 17 p.
2011-19	Eveline van Leeuwen Peter Nijkamp	Operational advances in tourism research, 25 p.
2011-20	Aliye Ahu Akgün Tüzin Baycan Peter Nijkamp	Creative capacity for sustainable development: A comparative analysis of European and Turkish rural regions, 18 p.
2011-21	Aliye Ahu Gülümser Tüzin Baycan-Levent Peter Nijkamp	Business dynamics as the source of counterurbanisation: An empirical analysis of Turkey, 18 p.
2011-22	Jessie Bakens Peter Nijkamp	Lessons from migration impact analysis, 19 p.
2011-23	Peter Nijkamp Galit Cohen-blankshtain	Opportunities and pitfalls of local e-democracy, 17 p.
2011-24	Maura Soekijad Irene Skovgaard Smith	The 'lean people' in hospital change: Identity work as social differentiation, 30 p.
2011-25	Evgenia Motchenkova Olgerd Rus	Research joint ventures and price collusion: Joint analysis of the impact of R&D subsidies and antitrust fines, 30 p.
2011-26	Karima Kourtit Peter Nijkamp	Strategic choice analysis by expert panels for migration impact assessment, 41 p.
2011-27	Farook Lazrak Peter Nijkamp Piet Rietveld Jan Rouwendal	The market value of listed heritage: An urban economic application of spatial hedonic pricing, 24 p.
2011-28	Peter Nijkamp	Socio-economic impacts of heterogeneity among foreign migrants: Research and policy challenges, 17 p.
2011-29	Masood Gheasi Peter Nijkamp	Migration, tourism and international trade: Evidence from the UK, 8 p.
2011-30	Karima Kourtit	Evaluation of cyber-tools in cultural tourism, 24 p.

	Peter Nijkamp Eveline van Leeuwen Frank Bruinsma	
2011-31	Cathy Macharis Peter Nijkamp	Possible bias in multi-actor multi-criteria transportation evaluation: Issues and solutions, 16 p.
2011-32	John Steenbruggen Maria Teresa Borzacchiello Peter Nijkamp Henk Scholten	The use of GSM data for transport safety management: An exploratory review, 29 p.
2011-33	John Steenbruggen Peter Nijkamp Jan M. Smits Michel Grothe	Traffic incident management: A common operational picture to support situational awareness of sustainable mobility, 36 p.
2011-34	Tüzin Baycan Peter Nijkamp	Students' interest in an entrepreneurial career in a multicultural society, 25 p.
2011-35	Adele Finco Deborah Bentivoglio Peter Nijkamp	Integrated evaluation of biofuel production options in agriculture: An exploration of sustainable policy scenarios, 16 p.
2011-36	Eric de Noronha Vaz Pedro Cabral Mário Caetano Peter Nijkamp Marco Painho	Urban heritage endangerment at the interface of future cities and past heritage: A spatial vulnerability assessment, 25 p.
2011-37	Maria Giaoutzi Anastasia Stratigea Eveline van Leeuwen Peter Nijkamp	Scenario analysis in foresight: AG2020, 23 p.
2011-38	Peter Nijkamp Patricia van Hemert	Knowledge infrastructure and regional growth, 12 p.
2011-39	Patricia van Hemert Enno Masurel Peter Nijkamp	The role of knowledge sources of SME's for innovation perception and regional innovation policy, 27 p.
2011-40	Eric de Noronha Vaz Marco Painho Peter Nijkamp	Impacts of environmental law and regulations on agricultural land-use change and urban pressure: The Algarve case, 18 p.
2011-41	Karima Kourtit Peter Nijkamp Steeff Lowik Frans van Vught Paul Vulto	From islands of innovation to creative hotspots, 26 p.
2011-42	Alina Todiras Peter Nijkamp Saidas Rafijevas	Innovative marketing strategies for national industrial flagships: Brand repositioning for accessing upscale markets, 27 p.

2011-43	Eric de Noronha Vaz Mário Caetano Peter Nijkamp	A multi-level spatial urban pressure analysis of the Giza Pyramid Plateau in Egypt, 18 p.
2011-44	Andrea Caragliu Chiara Del Bo Peter Nijkamp	A map of human capital in European cities, 36 p.
2011-45	Patrizia Lombardi Silvia Giordano Andrea Caragliu Chiara Del Bo Mark Deakin Peter Nijkamp Karima Kourtiti	An advanced triple-helix network model for smart cities performance, 22 p.
2011-46	Jessie Bakens Peter Nijkamp	Migrant heterogeneity and urban development: A conceptual analysis, 17 p.
2011-47	Irene Casas Maria Teresa Borzacchiello Biagio Ciuffo Peter Nijkamp	Short and long term effects of sustainable mobility policy: An exploratory case study, 20 p.
2011-48	Christian Bogmans	Can globalization outweigh free-riding? 27 p.
2011-49	Karim Abbas Bernd Heidergott Djamil Aïssani	A Taylor series expansion approach to the functional approximation of finite queues, 26 p.